

# ValidBraindumps

Over **61842+** Satisfied Customers

About Us

ValidBraindumps

HOME CERTIFICATIONS HOW TO PAY? GUARANTEE FAQ CART (0)

Test4engine

WE

TRY BEFORE YOU BUY

Download a free sample of any of our exam questions and answers

- > 24/7 customer support, Secure shopping site
- > Free One year updates to match real exam scenarios
- > If you failed your exam after buying our products we will refund the full amount back to you.

IGINE

st version

n exams. Besides for the

ear test questions and

to the highest standards of technical

fronter experts and published authors for

Select a vendor... Select an exam... Your email address Free Download

## Try Before You Buy

Download a free sample of any of our exam questions and answers

- > 24/7 customer support, Secure shopping site
- > Free One year updates to match real exam scenarios
- > If you failed your exam after buying our products we will refund the full amount back to you.

<http://www.validbraindumps.com>

Free valid test braindumps for IT certification valid exam

**Exam** : **CPR**

**Title** : Cardiopulmonary resuscitation

**Vendor** : ISQI

**Version** : DEMO

NO.1 What is generally the best way to learn how to use a defibrillator?

- A. By watching E.R.
- B. Buy an AED Trainer
- C. Learn as you go when the time comes
- D. Take a training class

**Answer:** D

NO.2 You are performing chest compressions for an adult victim of cardiac arrest. Which of the following is the correct rate of compressions for adult CPR?

- A. a rate of 150 times a minute
- B. a rate of 100 times a minute
- C. a rate of 80 times a minute
- D. a rate of 60 times a minute

**Answer:** B

NO.3 You have powered ON your AED and attached electrode pads and cables. What should you do NEXT?

- A. Call "Clear!" (check that no one is touching the victim); then: press ANALYZE
- B. Give 2 breaths and 15 chest compressions
- C. Press the SHOCK button
- D. Recheck for a pulse

**Answer:** A

Explanation:

The scenario puts the rescuer at the exact point in the sequence where everything is ready for rhythm assessment and shock (if indicated). The answer reminds the learner that you must always remember to "clear" before both analysis and shocks.

Answers b, c, and d are incorrect because they are each out of the correct sequence.

NO.4 Which statement is incorrect?

- A. Suctioning attempts should last 10 seconds or less.
- B. Suctioning should be followed by the administration of 100% oxygen.
- C. Yankauer catheters are used to suction the nose.
- D. Monitor the patient's heart rate and clinical appearance during suctioning

**Answer:** C

Explanation:

Yankauer catheters are used to suction the nose.

NO.5 When performing adult CPR, the depth of compression should be:

- A. at least 2 inches deep
- B. 2 1/2 inches deep
- C. 3 inches deep
- D. None of the above

**Answer:** A

NO.6 Which of the following statements best describes the effects of chest compressions and ventilations during adult CPR?

- A. they decrease the need for coronary artery bypass
- B. they shock the heart in VF(ventricular fibrillation) back to normal heart rhythm
- C. they constrict the pupils to help preserve vision
- D. they provide a flow of oxygen to the heart and brain

**Answer:** D

NO.7 A 7-year-old child is pulled from below the water of a neighbor's pool. The neighbor's children say the child did not fall or jump into the water - she was swimming and suddenly slipped under the water. You kneel beside the child and find that she is unresponsive. You send someone to phone 911 while you remain with the child. What should you do next?

- A. turn the child's head down and perform the Heimlich maneuver to drain water from the lungs
- B. give five back blows, then do a finger sweep of the mouth
- C. find the proper hand position and begin chest compressions
- D. open the airway with the head-tilt-chin-lift maneuver, look-listen-feel for breathing, and if the child is not breathing normally, give two rescue breaths

**Answer:** D

NO.8 CORRECT TEXT

What does AED stand for?

**Answer:** Automated External Defibrillator

NO.9 When you deliver the first rescue breath, what should you do if the victim's chest does not rise?

- A. Deliver a sharp blow to the center of the chest
- B. Give another breath
- C. Perform the head tilt-chin lift
- D. Stop performing CPR

**Answer:** C

NO.10 What is the recovery position?

- A. Placing the victim on his or her side
- B. When the victim stands up
- C. Raising the feet up above the heart
- D. Placing the victims in a sitting position

**Answer:** A

NO.11 If you find an unresponsive victim with no evidence of injury. how should you open the airway

- A. tilt the head and lift the chin
- B. turn the head to one side, then open the mouth
- C. perform abdominal thrusts (the Heimlich maneuver)
- D. wipe out the mouth with a gauze pad

**Answer:** A

NO.12 At what age do children become adults in CPR terms?

- A. 8
- B. 12
- C. 16
- D. 18

**Answer:** B

NO.13 Which of the following victims should NOT be treated with an AED?

- A. victim less than 8 years of age
- B. victim with a history of severe diabetes and a prosthetic (artificial leg)
- C. victim who is recovering from recent heart surgery (scar on chest)
- D. victim who is recovering from a recent stroke

**Answer:** A

Explanation:

At this time, AEDs are not approved for use for cardiac arrest victims under the age of 8.

Victims with a pacemaker, heart surgery, or recent stroke can safely be treated and shocked by an AED.

NO.14 When attempting defibrillation, how many shocks with an AED should be delivered before resuming CPR?

- A. 1
- B. 2
- C. 3
- D. 4

**Answer:** A

NO.15 If the victim is breathing normally, but not responding you should:

- A. Proceed with administering CPR
- B. Give mouth to mouth
- C. Turn the victim to the side and wait for emergency personnel
- D. None of the above

**Answer:** C